


Available online at <http://ajprd.com>

Asian Journal of Pharmaceutical Research and Development

(An International Peer-Reviewed Journal of Pharmaceutical Research and Development)

Open Access to Pharmaceutical and Medical Research

© 2013-18, publisher and licensee AJPRD, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open Access

Research Article

COMPARATIVE STUDY OF ANTIBACTERIAL PROPERTIES OF FLAVONOIDS OF LEAVES FROM DIFFERENT CACTUS, PERENNIAL GRASSES AND MEDICINAL PLANT

Singariya Premlata ^{*1}, Mourya Krishan Kumar² and Kumar Padma¹¹Laboratory of Tissue Culture and Secondary Metabolites, Department of Botany, University of Rajasthan, Jaipur-302004.²Animal Disease Diagnostic Laboratory, Rural Veterinary Polyclinic, Hingonia, Jaipur(Rajasthan).

ABSTRACT

Comparative study of antibacterial properties of flavonoids of leaves from different Cactus (*Euphorbia caducifolia*); perennial grasses (*Panicum antidotale* and *Lasiurus indicus*) and medicinal plant (*Anaegissus rotundifolia*) against some important bacteria (G+ve or G-ve) was done. Flavonoid extracts of leaves from all the selected plants were found to possess strong antibacterial activity against these test pathogenic microbes, as revealed by ZOI=(mm± SD), AI, MIC=(mg/ml), MMC=(mg/ml) and TA=(ml/gm) of extracts against each sensitive test pathogenic bacteria were also evaluated. Most of the pathogens were found to be sensitive against these flavonoid extracts. Free flavonoid extract of grasses and medicinal plant showed higher activity against most of the pathogens. *Euphorbia caducifolia* is the most inactive plant in this study. All pathogens didn't show any activity against *Euphorbia caducifolia* except *Proteus mirabilis* which show 7.83 ± 0.21 mm ZOI for free flavonoid. Highest ZOI (18.67 ± 0.26) was recorded for free flavonoid of medicinal plant (*Anaegissus rotundifolia*) against *Bacillus subtilis* (G+ve bacteria) as well as other parameters like AI (1.245), MIC (0.039 mg/ml), MMC (0.78 mg/ml) were found higher against *Bacillus subtilis* and highest TA (1012.821 ml/gm) were found highest for free flavonoid of *Lasiurus indicus* against *Bacillus subtilis*. The findings of the present study suggested the exploitation of flavonoid extracts of *Anaegissus rotundifolia* and of *Lasiurus indicus* for future antibacterial drugs.

Key words: Antibacterial, Flavonoids, *Lasiurus indicus*, *Anaegissus rotundifolia*, Zone of Inhibition, Activity Index, Minimum Bactericidal Concentration and Total Activity.



Cite this article as:

Singariya Premlata, Comparative Study of Antibacterial Properties of Flavonoids of Leaves from Different Cactus, Perennial Grasses and Medicinal Plant, Asian Journal of Pharmaceutical research and Development. 2018;6(3):32-41 DOI: <http://dx.doi.org/10.22270/ajprd.v6.i2.372>

*Address for Correspondence

Premlata Singariya, Laboratory of Tissue Culture and Secondary Metabolites, Department of Botany, University of Rajasthan, Jaipur, India

INTRODUCTION:

Many commercially proven drugs used in modern medicine were initially used in crude form in traditional or folk healing practices or for other purposes that suggested potentially useful biological activity (1). Antibiotics were medical miracles during the Second World War but are now becoming impotent bacterial weaponry (2). This has caused an urgent need for the research of new and innovative ways to control bacterial invasions especially by multi-resistant pathogens such as *B. subtilis* (Gram +ve) and *P. aeruginosa* (Gram -ve) (3).

Flavonoids are hydroxylated phenolic substances and are known to be synthesized by plants in response to microbial infection (4). Their activities are structure dependent. The chemical nature of flavonoids depends on their structural class, degree of hydroxylation, other substitutions and conjugations, and degree of polymerization. Recent interest in these substances has been stimulated by the potential health benefits arising from the antioxidant activities of the polyphenolic compounds. Functional hydroxyl groups in flavonoids mediate their antioxidant effects by scavenging free radicals and/or by chelating metal ions. These bioactive compounds exert antimicrobial activity through different

mechanisms. Natural antioxidants mainly come from plants in the form of phenolic compounds such as flavonoid, phenolic acids, tocopherols etc. (5). Phenolic compounds exhibit a considerable free radical scavenging (antioxidant) activity (6). The plants are vital source of innumerable number of antimicrobial compounds. Several phytochemical constituents like flavonoids (7), phenolics and polyphenols (8), tannins (9), etc. are effective antimicrobial substances against a wide range of microorganisms. One of the largest groups of chemical produced by plant is the alkaloids and their amazing effects on humans have led to the development of powerful painkiller medications (10).

Infectious diseases are the world's leading cause of premature deaths. In recent years, drug resistance to human pathogenic bacteria has commonly been reported from all over the world (11). Even though pharmaceutical companies produce number of new antibacterial drugs, but gradual resistance to these drugs has increased which is matter of global concern besides synthetic drugs are normally associated with side effects (hypersensitive, immune suppression etc). Use of phytochemicals with known antimicrobial properties can be of great significance in therapeutic treatments. Present study is an effort towards this direction. Results reveal that all the tested extracts of selected plant exhibited growth inhibitory activity against one of the other bacterial strains selected.

Panicum antidotale and *Lasiurus indicus* (perennial grasses) are more efficient at gathering Carbon dioxide and utilizing nitrogen from the atmosphere and recycled Nitrogen in the soil (12,13). Perennial grasses are more competitive under the conditions of high temperature, solar radiation and low moisture. These grasses have excellent soil binding capacity which helps to conserve soil in desert areas. These grasses are distributed in hotter and drier parts of India, Mediterranean region, tropical and southern Africa. This grass, fed green, turned into silage, or made into hay is said to increase flow of milk in cattle and impart a sleek and glossy appearance (14). Seeds of this grass are used as famine food by the tribal during severe conditions.

Klebsiella pneumonia more frequently causes lung destruction and pockets of pus in the lung (known as empyema), respiratory infections, such as bronchitis, which is usually a hospital-acquired infection (15). *Proteus mirabilis* cause obstruction and renal failure. It

can also cause wound infections, septicemia and pneumonias, mostly in hospitalized patients. *A. tumefaciens* (Plant pathogen) uses horizontal gene transfer to cause tumors "crown gall disease" in plants. It can be responsible for opportunistic infections in humans with weakened immune systems (16). *R. planticola* has been determined to cause severe pancreatitis in one case (17). *Bacillus subtilis* can contaminate food; however, they seldom result in food poisoning. *E. aerogens* is a nosocomial and pathogenic bacterium that causes opportunistic infections including most types of infections.

MATERIALS AND METHODS:

Experimental design: Flavonoid extract of leaves of selected plants cactus (*Euphorbia caducifolia*); perennial grasses (*Panicum antidotale* and *Lasiurus indicus*) and medicinal plant (*Anaegissus rotundifolia*) were prepared by hot extraction method (18) in Soxhlet assembly.

All the flavonoid extracts were then screened for evaluation of antibacterial activity by Disc Diffusion Assay (DDA) (19,20), in terms of Zone of Inhibition [ZOI=(mm± SD)], Activity Index (AI), Minimum inhibitory concentration [MIC=(mg/ml)], Minimum Bactericidal concentration [MBC=(mg/ml)] and Total activity [TA=(ml/gm)] of extracts against some medically important pathogenic microbes (*Bacillus subtilis*=G+ve, *Escherichia coli*, *Raoultella planticola*, *Enterobacter aerogens*, *Proteus mirabilis*, *Klebsiella pneumonia* and *Agrobacterium tumefaciens*=G-ve bacteria. The fraction showing best activity was then used for determining of minimum inhibitory concentration (MIC) by tube dilution method (21) and minimum bactericidal concentration (MBC).

Plant material: Leaves of selected plants *Euphorbia caducifolia*, *Panicum antidotale*, *Lasiurus indicus* and *Anaegissus rotundifolia* were collected in the month of August 2014 from the Central Arid Zone Research Institute (CAZRI), Jodhpur (Rajasthan). Plant samples were identified and deposited in the herbarium, Department of Botany, University of Rajasthan, Jaipur. Herbarium no. of these plants are recorded in table 1. The collected plant materials were transferred immediately to the laboratory cleaned with water (22) and selected plant parts were separately shade dried (23) until weight has been constant (24) (table-1).

Table 1: Selected plants:

S. No.	Name of plant	Herbarium No.
1.	<i>Euphorbia caducifolia</i>	RUBL211367
2.	<i>Panicum antidotale</i>	RUBL211364
3.	<i>Lasiurus indicus</i>	RUBL211366
4.	<i>Anaegissus rotundifolia</i>	RUBL211363

Table 2: Flavonoid Extraction by hot extraction method from leaves of selected plants (*Euphorbia caducifolia*, *Panicum antidotale*, *Lasiurus indicus* and *Anaegissus rotundifolia*):

S. No.	Name of plant	Flavonoid	
		Free Flavonoid	Bound Flavonoid
1.	<i>Euphorbia caducifolia</i>	+	+
2.	<i>Panicum antidotale</i>	+	+
3.	<i>Lasiurus indicus</i>	+	+
4.	<i>Anaegissus rotundifolia</i>	+	+

Preparation of plant extracts: The shade-dried leaves were powdered with the help of a grinder (25) and passed through 40mm meshes (26) and stored in clean container for further use (27). The dried powder material was extracted by hot extraction method (28) using the Soxhlet apparatus (29).

Determination of total Flavonoid Content: The total flavonoid content was determined according to the aluminum chloride colorimetric method (30). Rutin was chosen as a standard (the concentration range; 0.005 to 0.1mg mL⁻¹) and the total flavonoid content was expressed as milligram per gram of dry extracts.

Preliminary detection of flavonoids: Following methods were used to determine the presence of flavonoids in the plant samples:

Five 5ml of dilute ammonia solution was added to a portion of the aqueous filtrate of each plant extract, followed by addition of concentrated H₂SO₄. A yellow color observed in each extract indicated the presence of flavonoids. The yellow color disappeared on standing.

Ethyl acetate test: A portion of the powdered plant sample was heated with 10 ml of ethyl acetate over a steam bath for 3 min. The mixture was filtered and 4ml of the filtrate was shaken with 1ml of dilute ammonium solution. A yellow coloration was observed indicating a positive test for flavonoids.

Extraction of flavonoids: Collected plant leaves were separately shade dried, finely powered using a blender and subjected to extraction following the method of Subramanian and Nagarjan (29). About 100 g of each finely powered sample was Soxhlet extracted with 80% hot methanol (500 ml) on a water bath for 24 h and filtered. Each filtrate was re-extracted successively with petroleum ether (fraction I), ethyl ether (fraction II) and ethyl acetate (fraction III) using separating funnel. Petroleum ether fractions were discarded as being rich in fatty substances, whereas ethyl ether and ethyl acetate fractions were analysed for free and bound flavonoids,

respectively. Ethyl acetate fraction of each of the samples was hydrolysed by refluxing with 7% H₂SO₄ for 2 h (for removal of bounded sugars from the flavonoids) and filtered. The filtrate was extracted in ethyl acetate and washed with distilled water to neutrality. Ethyl ether (free flavonoid) and ethyl acetate fractions (bound flavonoids) thus obtained were dried in vacuo and weighed. The extracts were stored at 4°C and were re-suspended in their respective solvents to get 10 mg/ml concentration for antimicrobial assay.

Flavonoid extraction: Leaves of selected plants were taken for flavonoids extraction following the well-established method (29). Each extract was dried in vacuo and stored at 4 °C in airtight vials for further use. Percent (%) extractive values were calculated by the following formula-

Weight of dried extract

$$\text{Percent (\%) Extracts} = \frac{\text{Weight of dried extract}}{\text{Weight of dried plant material}} \times 100$$

Weight of dried plant material

Drugs and Chemicals Used:

Drugs: Streptomycin, Ciprofloxacin, Ceftriaxone (Standard Antibiotics for bacteria)

Chemicals: Sabouraud Dextrose Agar Medium (SDA) (Peptone 10g; Dextrose 20g; Agar 20g in 1000ml of distilled water; pH adjusted to 6.8-7.0 at 27±2°C), Ammonium chloride, ammonium solution, Ethyl acetate, H₂SO₄.

Micro-organisms: The organisms used in this study were seven bacteria (six G^{-ve} and one G⁺). Selected bacteria were procured from IMTECH, Chandigarh, India. Bacterial strains were grown and maintained on Muller-Hinton Agar Medium (MHA) at 37±2°C and sub cultured regularly (after every 30 days) and stored at 4°C as well as at -80°C by preparing suspensions in 10% glycerol (31) (table-3).

Table 3: Selected pathogens:

S. No.	Pathogens Type	Name of Pathogens	G ⁺ -ve/ G ^{-ve}	Specimen no.
1.	Bacteria	<i>Escherichia coli</i>	G ^{-ve}	MTCC-46
2.		<i>Raoultella planticola</i>	G ^{-ve}	MTCC-530
3.		<i>Bacillus subtilis</i>	G ⁺ -ve	MTCC-121
4.		<i>Enterobacter aerogens</i>	G ^{-ve}	MTCC-111
5.		<i>Proteus mirabilis</i>	G ^{-ve}	MTCC-3310
6.		<i>Klebsiella pneumoniae</i>	G ^{-ve}	MTCC-4030
7.		<i>Agrobacterium tumefaciens</i>	G ^{-ve}	MTCC-431

Preparation of Test Pathogens and Disc Diffusion Assay (DDA): Initial screening of different extracts for their antibacterial activity carried out using MHA and NA media but, did not reveal any significant difference, thus further studies were carried out using NA medium

only (32). Bacterial strains were grown and maintained on NA medium. Bacterial growth can observe after a minimum of 18 hours and occasionally until 24 hours. Disc diffusion assay (DDA) was performed for screening. NA base plates were seeded with bacteria

(inoculum size 1×10^8 CFU/ml for bacteria). Sterile filters paper discs (Whatman no. 1, 5mm in diameter) were impregnated with 100 μ l of each of the extracts (100 mg/ml) to give a final concentration of 1 mg/disc and left to dry in vacuo so as to remove residual solvent, which might interfere with the determination (33). Petri plates were pre-seeded with 15 ml of growth agar medium and 1.0 ml of inoculum (34,35). Extract discs were then placed on the preseeded agar plates. Each extract was tested with Streptomycin (10mcg/disc), Ciprofloxacin (10mcg/disc) and Ceftriaxone (10mcg/disc) commercial discs of antibiotics as positive control (standard) and experiment was done thrice (36) for bacteria. The plates were kept at 4°C for 1 h for diffusion of extract, thereafter were incubated at 37°C for bacteria (24 h) and 27°C (37). Zone of Inhibition (ZOI) or depressed growth of microorganisms was measured. This method was followed by various researchers (38,39).

Zones of Inhibition (ZOI): Zones of Inhibition [ZOI=(mm \pm SD)], measured in mm (mean value; include 5 mm diameter of disc), indicate that no bacterial growth around the tested flavonoid extract disc. ZOI measured and compared with the standard reference antibiotics (40-42).

Activity index (AI): Activity index for each extract was calculated by the formula (43).

Inhibition Zone of the sample

Activity index (AI) = -----

Inhibition Zone of the standard

Determination of Minimum Inhibitory Concentration (MIC) by Serial Dilution Method/ Micro Broth Dilution (MBD) Method: Minimum inhibitory concentrations (MICs) are considered as the "gold standard" for determining the susceptibility of the organisms to antimicrobials. MIC was determined for plant extracts those showing antimicrobial activities against test pathogens in disc diffusion assay (44). Broth micro-dilution method (45) was followed for determination of MIC values. Plant extracts were re-suspended in acetone (which has no activity against test microorganisms) to make 10mg/ml final concentration and then was added to test tubes containing 1 ml of sterile NA media. The tubes were then inoculated with a drop of microbial suspension (for bacteria 1×10^8 CFU/ml) and the tubes were incubated at $37 \pm 2^\circ\text{C}$ for 24 hrs in a BOD (Biological Oxygen Demand) incubator. Each extract was assayed in duplicate and each time two sets of microtiter plates were prepared, one was kept for incubation while another set was kept at 4°C for comparing the turbidity (46) in the wells of microtiter plate. A tube containing nutrient broth without extract was taken as control. The least extra concentration which inhibited the growth of the test organisms was taken as MIC (47). Broth media of 96-wells of microtiter plates using two serial dilutions. There after 100 μ l inoculum of standard size was added to each well. Bacterial and fungal suspensions were used as negative control, while broth containing standard drug was used as positive control. The MIC values were taken as the lowest concentration of the extracts in the well of the

microtiter plate that showed no turbidity after incubation. The turbidity of the wells in the microtiter plate was interpreted as visible growth of bacteria.

The MIC values were taken as the lowest concentration of the extracts in the test tubes that showed no turbidity after incubation (48). The turbidity of the test tube was interpreted as visible growth of microorganisms.

Determination of Minimum bactericidal concentration (MBC) by Serial Dilution Method/ Micro Broth Dilution (MBD) method: It is defined as

the concentration of the antimicrobial that results in a 99.9% reduction in CFU/ml for bacteria, compared with the organism concentration in the original inoculum (49). Equal volume of various concentration of each extract and nutrient agar were mixed in micro-tubes to make up 0.5ml of solution. Then 0.5ml of McFarland standard of the organism suspension was added to each tube (50). The tubes were incubated aerobically at 37 °C for 24 h for bacteria in a BOD incubator and observed for change in turbidity after 24 hrs and 48 hrs comparisons with the growth and sterility controls. Two control tubes were maintained for each test batch. These include tube-containing extract without inoculum and the tube containing the growth medium and inoculum. The MBC was determined by sub culturing the test dilution on Nutrient Agar followed by incubation. The MBC was determined by sub-culturing 50 μ l from each well showing no apparent growth. Least concentration of extract showing no visible growth on sub-culturing was taken as MBC (51). MBC was calculated for those extracts that had shown high antimicrobial activity against tested organisms.

Total activity (TA-mg/ml): Total activity is the volume at which test extract can be diluted without losing the ability to kill microorganisms (52). It is calculated by dividing the amount of extract from 1 g plant material by the MIC of the same extract or compound isolated and is expressed in ml/gm. In mathematical terms it can be expressed as:

Amount extracted from 1g plant material

Total activity (TA) = -----

MIC of the extract

Statistical Analysis: Mean value and standard deviation were calculated for each test pathogens. Data were analyzed by one-way ANOVA and p values ($p > 0.005$) were considered significant (53).

RESULTS:

Preliminary phyto-profiling: The preliminary phyto-profiling of flavonoid extraction from leaves of selected plants were carried out (table 3) according to Farnsworth (54). Free flavonoid show higher yield as compare to bound flavonoid of all the selected plants. The highest yield (mg/gm w/w in dry weight) was recorded (table 4) for free flavonoid of *L. indicus* (39.5) followed by *P. antidotale* (34.52).

Table 4: Preliminary phyto-profile of Flavonoid Extraction (Free Flavonoid and Bound Flavonoid) from leaves of selected plants:

S. No.	Phytochemical estimation	<i>Euphorbia caducifolia</i>		<i>Panicum antidotale</i>		<i>Lasiurus indicus</i>		<i>Anaegissus rotundifolia</i>	
		Free	Bound	Free	Bound	Free	Bound	Free	Bound
1.	% Yield	1.86	1.78	3.45	2.94	3.95	3.29	1.84	1.70
2.	Yield in mg/gm DW	18.6	17.8	34.52	29.38	39.5	32.9	18.43	6.143
3.	Yield in gm (30gm)	6.200	5.933	11.507	9.793	13.167	10.967	16.96	5.653
4.	Colour	Green	Green	Dark Green	Greenish Yellow	Brown	Greenish Yellow	Dark green	Green

Antibacterial activity: Extraction of flavonoid from leaves of selected plants were found to possess strong antibacterial activity against these test pathogens, in terms of Zone of Inhibition [ZOI=(mm± SD)] and Activity Index (AI), Minimum inhibitory concentration [MIC=(mg/ml)], Minimum bactericidal concentration [MBC=(mg/ml)] and Total activity [TA=(ml/gm)] of extracts against each sensitive test pathogens were also evaluated.

Zone of Inhibition [ZOI=(mm± SD)]:

Antibacterial activity: Most susceptible organism in this study was *B. subtilis* (G+ve bacteria) against which, most of the plant extracts showed higher zone of inhibition as compare to the other organisms (G-ve bacteria). Free flavonoid show higher activity as compare to bound flavonoid of all the selected plants. Highest ZOI were recorded for leaves of *Anaegissus rotundifolia* (18.67±0.26 mm) followed by *Lasiurus indicus* (18.50±0.64 mm) against *B. subtilis* (G+ve bacteria) (table-5).

Table 5: Zone of Inhibition (ZOI) (mm± SD) of Flavonoid extraction from leaves of selected plants (*Euphorbia caducifolia*, *Panicum antidotale*, *Lasiurus indicus* and *Anaegissus rotundifolia*) against tested pathogens (G+ve and G-ve bacteria):

S. No.	Name of Pathogen	<i>Euphorbia caducifolia</i>		<i>Panicum antidotale</i>		<i>Lasiurus indicus</i>		<i>Anaegissus rotundifolia</i>	
		Free	Bound	Free	Bound	Free	Bound	Free	Bound
1.	<i>Escherichia coli</i>	-	-	16.67± 0.21	-	18.33± 0.24	-	16.50± 0.27	15.17± 0.67
2.	<i>Raoultella planticola</i>	-	-	8.50± 0.21	7.33± 0.26	9.33± 0.26	8.50± 0.64	12.33± 0.24	10.83± 0.26
3.	<i>Bacillus subtilis</i>	-	-	15.33± 0.26	-	18.50± 0.64	-	18.67± 0.26	16.83± 0.64
4.	<i>Enterobacter aerogens</i>	-	-	7.83± 0.26	7.33± 0.21	8.67± 0.24	7.33± 0.26	10.67± 0.24	9.33± 0.26
5.	<i>Proteus mirabilis</i>	7.83± 0.21	-	9.83± 0.26	7.50± 0.24	11.83± 0.24	8.67± 0.21	11.33± 0.24	9.50± 0.26
6.	<i>Klebsiella pneumoniae</i>	-	-	9.83± 0.26	7.33± 0.21	11.50± 0.64	7.33± 0.24	14.50± 0.21	11.67± 0.27
7.	<i>Agrobacterium tumefaciens</i>	-	-	8.67± 0.24	7.50± 0.26	9.83± 0.21	8.67± 0.24	12.33± 0.27	10.50± 0.24

Activity Index (AI): Most susceptible organism in this study was *B. subtilis* (G+ve bacteria) against which, most of the plant extracts showed higher activity as compare to the other organisms (G-ve bacteria). Free

flavonoid show higher activity as compare to bound flavonoid of all the selected plants. Highest AI values were recorded for *A. rotundifolia* (1.245) followed by *L. indicus* (1.233) against *B. subtilis* (table-6).

Table 6: Activity index (AI) of Flavonoid extraction from leaves of selected plants (*Euphorbia caducifolia*, *Panicum antidotale*, *Lasiurus indicus* and *Anaegissus rotundifolia*) against tested pathogens (G-ve and G+ve bacteria):

S. No.	Name of Pathogens	<i>Euphorbia caducifolia</i>		<i>Panicum antidotale</i>		<i>Lasiurus indicus</i>		<i>Anaegissus rotundifolia</i>	
		Free	Bound	Free	Bound	Free	Bound	Free	Bound
1.	<i>E. coli</i>	--	--	1.042	--	1.146	--	1.031	0.948
2.	<i>R. planticola</i>	--	--	0.567	0.489	0.622	0.567	0.822	0.722
3.	<i>B. subtilis</i>	--	--	1.022	--	1.233	--	1.245	1.122
4.	<i>E. aerogens</i>	--	--	0.489	0.458	0.542	0.458	0.667	0.583
5.	<i>P. mirabilis</i>	0.489	--	0.614	0.469	0.739	0.542	0.708	0.794
6.	<i>K. pneumoniae</i>	--	--	0.655	0.489	0.767	0.489	0.967	0.778
7.	<i>A. tumefaciens</i>	--	--	0.578	0.500	0.655	0.578	0.822	0.700

Minimum inhibitory concentration [MIC=(mg/ml)]: MICs were evaluated for those plant parts extracts, which had shown activity in 'Disc Diffusion assay'. The range of MIC of Alkaloid extracts recorded was 0.020 - 10 mg/ml. In the present investigation lowest MIC

values (0.039 mg/ml) were recorded for free flavonoid extracts of *Anaegissus rotundifolia* and *Lasiurus indicus* followed by bound flavonoid of *Anaegissus rotundifolia* (0.078 mg/ml) against *B. subtilis* (table 7).

Table 7: Minimum Inhibitory Concentration (MIC-mg/ml) of Flavonoid extraction from leaves of selected plants (*Euphorbia caducifolia*, *Panicum antidotale*, *Lasiurus indicus* and *Anaegissus rotundifolia*) against tested pathogens (G-ve and G+ve bacteria):

S. No.	Name of Pathogens	<i>Euphorbia caducifolia</i>		<i>Panicum antidotale</i>		<i>Lasiurus indicus</i>		<i>Anaegissus rotundifolia</i>	
		Free	Bound	Free	Bound	Free	Bound	Free	Bound
1.	<i>E. coli</i>	--	--	0.156	--	0.078	--	0.156	0.313
2.	<i>R. planticola</i>	--	--	2.500	5.000	2.500	2.500	2.500	1.250
3.	<i>B. subtilis</i>	--	--	0.156	--	0.039	--	0.039	0.078
4.	<i>E. aerogens</i>	--	--	5.000	5.000	5.000	5.000	1.250	2.500
5.	<i>P. mirabilis</i>	5.000	--	2.500	5.000	1.250	5.000	1.250	2.500
6.	<i>K. pneumoniae</i>	--	--	1.250	5.000	2.500	5.000	0.156	2.500
7.	<i>A. tumefaciens</i>	--	--	2.500	5.000	1.250	2.500	2.500	1.250

Minimum bactericidal Concentration (MBC = mg/ml): The highest dilution that yielded no single pathogen was taken as the Minimum microcidal concentration. The range of MBC of flavonoid extracts recorded was 0.020 - 10 mg/ml. In the present

investigation lowest MIC values (0.078mg/ml) were recorded for free flavonoid extracts of *Anaegissus rotundifolia* and *Lasiurus indicus* followed by bound flavonoid of *Anaegissus rotundifolia* (0.156 mg/ml) against *B. subtilis* (table 8).

Table 8: Minimum Bactericidal Concentration (MBC-mg/ml) of Flavonoid extraction from leaves of selected plants (*Euphorbia caducifolia*, *Panicum antidotale*, *Lasiurus indicus* and *Anaegissus rotundifolia*) against tested pathogens (G+ve and G-ve bacteria):

S. No.	Name of Pathogens	<i>Euphorbia caducifolia</i>		<i>Panicum antidotale</i>		<i>Lasiurus indicus</i>		<i>Anaegissus rotundifolia</i>	
		Free	Bound	Free	Bound	Free	Bound	Free	Bound
1.	<i>E. coli</i>	--	--	0.313	--	0.156	--	0.313	0.625
2.	<i>R. planticola</i>	--	--	2.500	5.000	2.500	5.000	5.000	1.250
3.	<i>B. subtilis</i>	--	--	0.313	--	0.078	--	0.078	0.156
4.	<i>E. aerogens</i>	--	--	10.000	10.000	10.000	10.000	2.500	5.000
5.	<i>P. mirabilis</i>	5.000	--	5.000	5.000	2.500	10.000	2.500	5.000
6.	<i>K. pneumoniae</i>	--	--	2.500	5.000	5.000	10.000	0.313	5.000
7.	<i>A. tumefaciens</i>	--	--	5.000	10.000	2.500	2.500	5.000	5.000

Total activity [TA=(ml/gm)]: Total activity indicated the volume upto which the extract could be diluted without losing the ability to kill the microorganisms. In the present investigation higher TA values were recorded for free flavonoids as compare to the bound flavonoids. Highest TA value was observed (1012.821

ml/gm) by free flavonoid extract against *Bacillus subtilis* followed by bound flavonoid extract (506.410 ml/gm) against *Escherichia coli* of *Lasiurus indicus* (table 9). It indicates that *Lasiurus indicus* has higher activity as compare to other selected plants.

Table 9: Total Activity (TA-ml/gm) of Flavonoid extraction from leaves of selected plants (*Euphorbia caducifolia*, *Panicum antidotale*, *Lasiurus indicus* and *Anaegissus rotundifolia*) against tested pathogens (G+ve and G-ve bacteria):

S. No.	Name of Pathogens	<i>Euphorbia caducifolia</i>		<i>Panicum antidotale</i>		<i>Lasiurus indicus</i>		<i>Anaegissus rotundifolia</i>	
		Free	Bound	Free	Bound	Free	Bound	Free	Bound
1.	<i>E. coli</i>	--	--	110.288	--	506.410	--	118.141	54.185
2.	<i>R. planticola</i>	--	--	6.904	11.752	15.800	13.160	7.372	13.568
3.	<i>B. subtilis</i>	--	--	221.282	--	1012.821	--	472.564	217.436
4.	<i>E. aerogens</i>	--	--	3.452	2.938	7.900	6.580	14.744	6.784
5.	<i>P. mirabilis</i>	3.72	--	6.904	5.876	31.600	6.580	14.744	6.784
6.	<i>K. pneumoniae</i>	--	--	6.904	11.752	15.800	6.580	118.141	6.784
7.	<i>A. tumefaciens</i>	--	--	6.904	23.504	31.600	13.160	7.372	13.568

DISCUSSION:

Most susceptible organisms observed in the investigations are *B. subtilis* and *Escherichia coli* against which, all the plant extracts showed good antibacterial activities. Highest antibacterial activity recorded by *Lasiurus indicus* against *B. subtilis* (G+ve bacteria).

Phytochemical constituents such as tannins, flavonoids, steroids and several other aromatic compounds of plant that serve as defense mechanisms against predation by many microorganisms, insects and herbivores. This may therefore explain the demonstration of antibacterial activity by the plant extracts. The demonstration of the antibacterial activity against bacteria may be indicative of the presence of broad spectrum antibiotic compounds. This will be of immense advantage in fighting the

menace of antibiotic refractive pathogens that are so prevalent in recent times.

G+ve bacteria *B. subtilis* was the most susceptible organism as compare to the G-ve bacteria (*E. coli*, *R. planticola*, *E. aerogens*, *P. mirabilis*, *K. pneumoniae* and *A. tumefaciens*) which supported the finding that plant extracts are usually more active against G+ve bacteria than G-ve bacteria (2,3,25,55-74). Susceptibility differences between G+ve and G-ve bacteria may be due to cell wall structural differences between these classes of bacteria. In the G-ve bacterial cell wall outer membrane appears to act as a barrier to many substances including synthetic and natural antibiotics (72).

Extracts under this study not only inhibit the bacterial growth but the ZOI developed, was more or less permanent when compared with the ZOI developed by

the standard drug used, as after sometime bacterial colonies could be easily seen in ZOI developed by standard drugs. In the light of the fact that micro-organisms are becoming resistant against the drugs in use, present investigation is of great significance, as far as the future drugs are concerned and the selected plants could be used by the pharmaceutical industries for preparing plant based antimicrobials drugs. Perennial grasses (*P. antidotale* and *L. indicus*) and *A. rotundifolia* easily grow in harsh climatic conditions or xeric conditions and require less care; hence its use as raw material for preparing drugs would definitely be economical.

Screening of all the selected plants (*Euphorbia caducifolia*, *Panicum antidotale*, *Lasiurus indicus* and *Anaegissus rotundifolia*) under investigation so far has not been worked out for flavonoids. Mostly crude extracts have been screened and that too without MIC,

MBC and TA determination. Such studies could only indicate their antibacterial potential but are not helpful in establishing them as an alternative for antibiotic. In the present study IZ, AI, MIC, MBC and TA have been evaluated for each extract. For most of the extracts MIC values recorded were very low, indicating strong bio-efficacy of the plant. In an overview of the bioactivity data obtained from the current investigation, it can be highlighted that the tested extracts have great potential to inhibit bacteria. There is a need for further investigation to explore the promising antibacterial properties of the selected plants.

ACKNOWLEDGEMENT:

Authors are thankful to the 'Head' Department of Botany, University of Rajasthan for providing all necessary facilities and to UGC for providing financial assistance for the present work.

REFERENCE:

- Retnam K, De-Britto A. Antimicrobial activity of a medicinal plant *Hybanthus enneaspermus* (Linn) F. Muell. Natural Product Rad. 2007;6(5): 366-368.
- Singariya P, Mourya KK, Kumar P. Estimation of Antibacterial Efficacy in Alkaloids of *Anaegissus rotundifolia* an Indigenous Medicinal Plant against Some Pathogenic Micro-organisms. Asian Journal of Research Chemistry. 2018a;11(2): 432-440.
- Lewis R. FDA Consumer Magazine. Retrieved from, 1995.
- Dixon RA, Dey PM, Lamb CJ. Phytoalexins: enzymology and molecular biology. Advances in Enzymology and Related Areas of Molecular Biology. 1983; 55:1-136.
- Ali SS, Kasoju N, Luthra A, Singh A, Sharanabasava H, Sahuand A, Bora U. Indian medicinal herbs as source of antioxidants. Food Res. Int., 2008; 41: 1-15.
- Wojdylo A, Oszmianski J, Czemerys R. Antioxidant activity and phenolic compounds in 32 selected herbs. Food Chemistry. 2007; 105: 940-949
- Tsuchiya H, Sato M, Miyazaki T, Fujiwara S, Tanigaki S, Ohyma M, Tanaka T, Ilinuma M. Comparative study on the antibacterial of phytochemical flavanones against methicillin resistant *Staphylococcus aureus*. Journal of Ethnopharmacology. 1996; 50: 27-34.
- Mason TL, Wassermann BP. Inactivation of red beet β -glucan synthase by inactive and oxidized phenolic compounds. Phytochemistry. 1987; 26: 2197-2202.
- Ya C, Gaffney SH, Lilley TH, Haslam E. Carbohydrate-polypheol complexation. 1988; 553. In: Hemingway, R.W. and Karches, J.J. (ed.), Chemistry and significance of condensed tannins. Plenum Press, New York.
- Raffauf RF. A Guide to Their Discovery and Distribution, Haworth Press Inc, New York, 1996;35.
- Piddock LJV, Wise R. Mechanisms of resistance to quinolones and clinical perspectives. Journal of Antimicrobial Chemotherapy. 1989; 23(4): 475-480.
- Singariya P. Effect of Sub-Optimal Environment and PGR's on Metabolic Pattern of Certain Species of *Cenchrus*. Ph.D Thesis, J. N. Vyas University, 2009.
- Bessman SP. Ammonia Metabolism in Animals: Symposium on Inorganic Nitrogen Metabolism. Mc Elroy and Glass (eds.) The Johns Hopkins Press 1956.
- Singariya P, Mourya KK, Kumar P. Evaluations of Organic Compounds in Ethyl Acetate extracts of Marwar Dhaman By gas chromatography- mass spectrometry. Journal of Plant Science. 33(1): 17-27.
- Martin WJ, PKW, Yu JA, Washington. Epidemiological significance of *Klebsiella pneumoniae* - a 3 month study. Mayo C. in. Proc, 1971; 46: 785-793.
- Dunne WM, Tillman J, Murray JC. Recovery of a strain of *Agrobacterium radiobacter* with a mucoid phenotype from an immuno compromised-child with bacteraemia. Journal of Clinical Microbiology. 1993; 31(9): 2541-2543. PMC 265809 PMID 8408587z
- Alves MS, Riley LW, Moreira BM. A case of severe pancreatitis complicated by *Raoultella planticola* infection. Journal of Medical Microbiology. 2007; 56(5): 696-698.
- Singariya P, Kumar P, Mourya KK. An activity guided isolation and evaluation of various solvent extracts of the leaves of Anjan grass. Hygeia journal for drugs and medicines. 2012s; 4(2): 49-56.
- Singariya P, Kumar P, Mourya KK. Evolution of Antibacterial Activity and Preliminary Phytochemical studies on the stem of *Cenchrus ciliaris* and *Cenchrus setigerus*. Asian Journal of Pharmaceutical and Clinical Research. 2012b; 5(1): 163-167.
- Bawer AW, Kirby WMM, Sherris JC, Turck M. Antibiotic susceptibility testing by a standardized single disc method. American Journal of clinical pathology. 1966; 45: 493-496.
- Singariya P, Kumar P, Mourya KK. Isolation of New Steroids and Evaluation of Bio-activity of Kala Dhaman Grass (*Cenchrus setigerus*). Brazilian Archives of Biology and Technology. 2014; 57(1): 62-69.
- Singariya P, Mourya KK, Kumar P. Identification of some bio-active compounds of iso-propyl alcohol extract of motha dhaman grass by gas chromatography-mass spectrometric analysis. Life Sciences Leaflets. 2016a; 72(2): 122-135.
- Singariya P, Kumar P, Mourya KK. Absence of Antibiotic Activities of *Cenchrus setigerus* and *Cenchrus ciliaris* Seed extracts in Different Polar Solvents. Journal of Pharmaceutical Negative Results. 2013b; 4(1): 71-75.
- Singariya P, Kumar P, Mourya KK. Evaluation of Antimicrobial Activity of Leaf extracts of Winter Cheery (*Withania somnifera*). International Journal of Pharm Tech Research. 2012p; 4(3): 1247-1253.
- Singariya P, Mourya KK, Kumar P. In-vitro Studies of Antimicrobial Activity of Crude Extracts of the Indian grasses Dhaman (*Cenchrus ciliaris*) and Kala-dhaman (*C. setigerus*). Indian Journal of Pharmaceutical Science. 2012w; 74(3): 261-265.
- Singariya P, Mourya KK, Kumar P. Gas Chromatography-Mass Spectrometric Analyses of Acetone extract of Marwar Dhaman grass for bio-active compounds. Plant Archives. 2015a; 15(2): 1065-1074.
- Singariya P, Kumar P, Mourya KK. Insignificant Antimicrobial Activities and Phyto-chemical screening in different extracts of Indian Ginseng. Journal of Pharmaceutical Negative Results. 2012o; 3(1): 41-45.

28. Singariya P, Mourya KK, Kumar P. Gas Chromatography-Mass Spectrometric analysis of acetone extract of *Cenchrus ciliaris* (Dhaman grass). International Journal of Science and Nature. 2015b; 6(4): 652-661.
29. Subramanian SS, Nagarajan S: Flavonoids of the seeds of *Crotalaria retusa* and *Crotalaria striata*. Current Science. 1969; 38: 65.
30. Lin JY, Tang CY. Determination of total phenolic and flavonoid contents in selected fruits and vegetables, as well as their stimulatory effects on mouse splenocyte proliferation. Food Chemistry. 2007; 101: 140-147.
31. Singariya P, Kumar P, Mourya KK. Bio-efficacy of Unripen Fruits of Winter Cherry (*Withania somnifera*). Research Journal of Pharmaceutical, Biological and Chemical Sciences. 2012f; 3(2): 479-487.
32. Grover J.K, Yadav S, Vats V. Medicinal plants of India with anti-diabetic potential. Journal of Ethnopharmacology. 2002; 81:81-100.
33. Singariya P, Kumar P, Mourya KK. Ripen Fruits of Indian Ginseng: Phyto-chemical and Pharmacological examination against Human and Plant Pathogens. International Journal of Applied Biology and Pharmaceutical Technology. 2012g; 3(2): 1-8.
34. Yamac M, Bilgili F. Antimicrobial activities of fruit bodies and/or mycelial cultures of some mushroom isolates. Pharmaceutical Biology. 2006; 44: 660-667.
35. Singariya P, Kumar P, Mourya KK. In-Vitro Antimicrobial Potency and Pharmacological Study of Flower extracts in different polar solvents of *Withania somnifera*. Asian Journal of Research Chemistry. 2012i; 5(3): 409-413.
36. Ravindra T, Russel H, Williams T. Antifungal activity of methanolic extract of *Capparis pepiaria* against the tested fungal strains. Journal of Ethnopharmacology. 2007; 8:109-114.
37. Aley-Kutty NA, Mathews SM, Leena PN. International Journal of Pharma and Bio Sciences. 2011; 2 (2) 182-187.
38. Sharma B, Kumar P. International Journal of Applied Research in Natural Products. 2009; 1(4) 5-12.
39. Kumar P, Sharma B, Bakshi N. Natural Product Research. 2009; 23(8): 719-723.
40. Singariya P, Kumar P, Mourya KK. Qualitative and Pharmacological examination of Root extracts of *Withania somnifera* against Human and Plant Pathogens. Asian Journal of Research in Chemistry. 2012m; 5(6): 733-737.
41. Harsh ML, Nag TN. Lloydia. 1984; 47, 365-368.
42. Cynthia H, O'Callaghan. Assessment of a new antibiotic. In: Hugo WB, Russel AD, editors. Pharmaceutical Microbiology. Oxford: Blackwell Scientific Publications, 1983, 122-134.
43. Singariya P, Kumar P, Mourya KK. Identification of New Bioactive Compounds by GC-MS and Estimation of Physiological and Biological Activity of Kala Dhaman (*Cenchrus setigerus*). International Journal of Pharmaceutical and Biological Archive. 2012r; 3(3): 610-616.
44. Joan SE. Clinical Bacteriology. 1975, Edward Arnold, London.
45. Barsi DF, Fan SH. The potential of aqueous and acetone extracts of galls of *Quercus infectoria* as antibacterial agents. Indian Journal of Pharmacology. 2005; 37(1): 26-29.
46. Demarsh P L, Gagnon RC, Hetzberg RP, Jaworski DD. Methods of Screening for antimicrobial compounds. Smithkline Beecham Corporation. Pub. World Intellectual Property Organization (WIPO) 2001.
47. Singariya P, Kumar P, Mourya KK. Isolation of Some New Steroids and Evaluation of Bio-activity of *Cenchrus ciliaris*. International Journal of Research in Pharmaceutical Science. 2012t; 3(4): 678-684.
48. Pepeljnjak S, Kalodera Z, Zovko M. Investigation of antimicrobial activity of *Pelarogarium radula* (Cav.) L'Herit. Acta Pharm. 2005; 55: 409-415.
49. Forbes BA, Sahm DF, Weissfeld AS. Bailey & Scott's Diagnostic Microbiology. 10th Ed. 2007, Mosby, Inc. Elsevier: St. Louis, Missouri, USA, 205.
50. Shahidi-Bonjar GH. Evaluation of Antibacterial properties of Iranian Medicinal plants against *Micrococcus aureus*, *Serratia marcescens*, *Klebsiella pneumoniae* and *Bordetella bronchiseptica*. Asian Journal of Science. 2004; 3(1): 82-86.
51. Delahaye C, Rainford L, Nicholson A, Mitchell S, Lindo J, Ahmad M. Antibacterial and antifungal analysis of crude extracts from the leaves of *Callistemon viminalis*. Journal of Medical and biological science. 2009; 3(1): 1-7.
52. Singariya P, Kumar P, Mourya KK. Comparative Bio-activity of Dhaman grass Root in Different Polar Solvents against Plant and Human Pathogens. International Journal of Green Pharmacy. 2012u; 6(3): 648-652.
53. Jain T, Sharma K. Assay of antibacterial activity of *Polyalthia longifolia* Benth. And Hook. Leaf extracts. Journal of Cell and Tissue Research. 2009; 9(2): 1817-1820.
54. Farnsworth NR. Biology and phytochemical screening of plants. Pharm Science. 1966; 55:225-276.
55. Singariya P, Mourya KK, Kumar P. Phytochemical and Antimicrobial Studies of Crude extract of Cow Sandbur and African Foxtail Grasses of Thar Desert. International Journal of Life Science. 2017a; 6(1): 1-10. DOI: 10.5958/2319-1198.2017.00001.X
56. Singariya P, Mourya KK, Kumar P. Phyto-profiling and antibiotic sensitivity test of ripe and unripe winter cherry. Journal of Pharmacy Research. 2017c; 11(11): 1376-1381.
57. Singariya P, Mourya KK, Gadi BR. Evaluation of Microcidal and Nitrogen assimilatory enzymes activity and identification of β - Sitosterol in C4 Grasses of Thar Desert. Environmental Impact on Biodiversity, 113-131. Editor: B. R. Bamniya and B. R. Gadi. Today & Tomorrow's Printers and Publishers, New Delhi - 110 002. 2016d; ISBN-p- 81-7019-547-7
58. Singariya P, Kumar P, Mourya KK. Antimicrobial activity and identification of 4,22-stigmastadiene -3-one and some other compounds in Motha Dhaman grass from Tribal area of Western Rajasthan. Proceedings of the National Academy of Sciences, India Section B: Biological Sciences. 2013a; 83(3):415-421. DOI 10.1007/s40011-012-0135-9.
59. Singariya P, Mourya KK, Kumar P. Phyto-chemical Screening and Antimicrobial Activities of Dhaman grass and Indian Ginseng. Journal of Pharmacy Research. 2012a; 5(1): 135-139.
60. Singariya P, Mourya KK, Kumar P. Antimicrobial Activity of the Crude Extracts of *Withania somnifera* and *Cenchrus setigerus* In-vitro. Pharmacognosy Journal. 2012d; 4(27): 60-65.
61. Singariya P, Kumar P, Mourya KK. Comparative Primary Phyto-profile and Microcidal Activity of *Cenchrus ciliaris* (Anjan grass) and *Withania somnifera* (Winter cherry). International Journal of Research in Ayurveda and Pharmacy. 2012e; 39(2): 303-308.
62. Singariya P, Kumar P, Mourya KK. Comparative Pharmacological Study of Fruits and Flowers Extract of *Withania somnifera*. International Journal of Pharmaceutical and Phyto-pharmacological Research. 2012h; 1(5): 276-282.
63. Singariya P, Kumar P, Mourya KK. Antibacterial and antifungal potential of some polar solvent extracts of Ashwagandha (Solanaceae) against the nosocomial pathogens. International Journal of Green Pharmacy. 2012j; 6(5): 17-22.
64. Singariya P, Kumar P, Mourya KK. Evolution of Indian Ginseng against Different Bacteria and Fungi. Asian

- Journal of Pharmaceutical and clinical Research. 2012k; 5(2): 145-148.
65. Singariya P, Kumar P, Mourya KK. Estimation of Bio-activity of Aerial parts of *Withania somnifera* Against the Bacterial and Fungal Microbes. International Journal of Pharmacy and Pharmaceutical science. 2012L; 4(3): 553-557.
66. Singariya P, Kumar P, Mourya KK. Screening for Antimicrobial Potency of Methanolic Extract of Indian Ginseng. International Journal of Pharmacy and Technology. 2012n; 4(2): 4537-4548.
67. Singariya P, Kumar P, Mourya KK. Identification of Steroid Compound using Preparative Thin Layer Chromatography, GC-MS & Anti-microbial and Antioxidant Properties of *Cenchrus setigerus* (Poaceae). International Journal of Pharmacy and Life Sciences. 2012q; 3(8): 1909-1916.
68. Singariya P, Kumar P, Mourya KK. *In-vitro* Bio-efficacy of Stem extracts of Ashwagandha against Some Pathogens. Journal of Current Pharmaceutical Research. 2011d; 8(1): 25-30.
69. Singariya P, Mourya KK, Kumar P. Comparative Microcidal Activity of *Withania somnifera* and *Cenchrus setigerus* against the Pathogenic Micro-organisms. International Journal of Pharmacy and Pharmaceutical science. 2011c; 3(5): 511-515.
70. Singariya P, Mourya KK, Kumar P. Bio activity of Crude Extracts of Leaves of *Cenchrus* Grass in different polar solvents against some Pathogenic Microbes. International Journal of Pharmaceutical Science Review and Research. 2011b; 11(1): 124-129.
71. Singariya P, Mourya KK, Kumar P. Preliminary Phyto-profile and Pharmacological Evaluation of some Extracts of *Cenchrus* grass against Selected Pathogens. Journal of Pharmaceutical Science and Research. 2011a; 3(8): 1387-1393.
72. Tortora GJ, Funke BR, Case CL. Microbiology: An Introduction, Benjamin Cummings, 2001. San Francisco.
73. Palombo EA, Semple SJ. Antibacterial activity of traditional Australian medicinal plants. Journal of Ethnopharmacology. 2001; 77: 151-157.
74. Lin J, Opake AR, Geheeb-Keller M, Hutchings AD, Terblanche SE, Jäger A K. Journal of Ethnopharmacology. 1999; 68: 267-274.