


Available online on 15.04.2020 at <http://ajprd.com>

## Asian Journal of Pharmaceutical Research and Development

Open Access to Pharmaceutical and Medical Research

© 2013-20, publisher and licensee AJPRD, This is an Open Access article which permits unrestricted non-commercial use, provided the original work is properly cited



Open Access

Research Article

## *Syzygium Polyanthum* Wight Leaf Extract Evaluation On *Aedes Spp* Instar III-IV Larvae

Widyawati Tri<sup>1,\*</sup>, Rizky Ilham<sup>2</sup><sup>1</sup>Department of Pharmacology and Therapeutic, Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia<sup>2</sup>Departement of Tropical Medicine, Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia

### ABSTRACT

*Aedes spp* mosquitos are the vectors that most cause diseases. The rise of insecticide resistance is related to the increasing of vectorial capacity. Plant can be used as an alternative source for controlling these vectors, one of which is *Syzygium polyanthum* (Wight). The present study was conducted to investigate the larvasidal properties of *S. polyanthum* leaf on *Aedes spp* instar III-IV. Ethanolic extract of *S. polyanthum* (EESP) was prepared by maceration using ethanol 70%. Larvae were divided into 7 groups (n=25, respectively). Group I to V consisted of different concentrations of EESP (100 (CI);150 (CII);200 (CIII);250 (CIV);300ppm (CV)); VI: Water (W) and VII : Temephos 1% (T). Larvicidal activity was evaluated by calculating the dead larva at 180, 360, 1440 and 2880 minutes to obtain Lethal Concentration 50% (LC50) and Lethal Time 50% (LT50) using Statistical Product and Service Solution (SPSS). The result showed that no larva found dead in W-treated group. Otherwise, the mortality of larva was increased with increasing of EESP concentration (CI to CV). LC50 and LT50 showed 213 ppm and 2410 minutes, respectively. We conclude that EESP has larvicidal activity on *Aedes spp* instar III-IV.

**Keywords:** *Aedes spp*, extract, larvicidal, *Syzygium polyanthum*

**ARTICLE INFO:** Received 16 Jan.2020; Review Completed 14 March 2020 ; Accepted 17 March 2020; Available online 15 April . 2020



Cite this article as:

Widyawati T, Rizky I, *Syzygium Polyanthum* Wight Leaf Extract Evaluation On *Aedes Spp* Instar Iii-Iv Larvae, Asian Journal of Pharmaceutical Research and Development. 2020; 8(2):07-09. DOI: <http://dx.doi.org/10.22270/ajprd.v8i2.678>

\*Address for Correspondence:

Tri Widyawati, Department of Pharmacology and Therapeutic, Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia

### INTRODUCTION

Dengue fever is the most important mosquito-borne viral disease of humans<sup>1</sup> which *Aedes spp* mosquitos act as the main vectors<sup>2</sup>. Annually, it is estimated 50-100 million cases with fatality rates between 0.5 and 3.5% in Asian countries<sup>3-5</sup>.

The chemical insecticides for controlling the mosquitos cycle are known to play a role in increasing mosquitos resistance<sup>6</sup>. Therefore, alternative source with larvicidal properties that obtained from plant could lead to the invention of new agents for vector control<sup>7</sup>. Plants that contained alkaloid, saponin, eugenol, flavonoids and tannin were reported able to kill *Aedes aegypti* larvae<sup>6,8</sup>. *Syzygium polyanthum* (*S. polyanthum*), a family of *Myrtaceae*, is widely used in Indonesia cuisines<sup>9</sup>. The potency of this plant as larvacid has been reported by Dwiyanti et al<sup>10</sup>. Their study showed that water extract of *S. polyanthum* had a killing power against *Aedes sp* larvae.

The bioactivity of plants extracts was related to their active compounds. Thus, the solvents that used to provide plant extracts also will affect the yield of its chemical compounds. The presents study was done to investigate the larvicidal activity of ethanolic extract of *S. polyanthum* (EESP) leaf on *Aedes spp* instar III-IV larvae.

### MATERIAL AND METHODS

The study was conducted on August-December 2018 at Pharmacology and Therapeutic Departement, Medical Faculty, Universitas Sumatera Utara, Medan, Indonesia.

#### *Extract preparation*

*S. polyanthum* leaves were obtained from Titi Kuning, Medan, North Sumatera, Indonesia. The fresh leaves were washed in running water and were dried in temperature room. The dried leaves or simplicia that had been grinded were extracted by maceration using ethanol 70% to obtain ethanol extract of *Syzygium polyanthum* (EESP).

### *Aedes spp* larvae

*Aedes spp* were obtained from Loka Litbang Kesehatan Pangandaran Ciamis, West Java, Indonesia (DP.02.01/1/831/2018).

### Experimental procedure

The larvae were divided into 7 groups (n=25) in 200 ml water (WHO, 2005) with 4 replications of each as follows: CI (EESP 100 ppm + Larvae); CII (EESP 150 ppm + Larvae); CIII (EESP 200 ppm + Larvae); CIV (EESP 250 ppm + Larvae); CV (EESP 300 ppm + Larvae); T (Temephos 1% + Larvae) served as positive control; W (Water + Larvae). Only the active *Aedes spp* larvae instar III-IV were included in the study.

### Data analysis

Data were analysed by Kruskal-Wallis and expressed as mean  $\pm$  SD. LC50 and LT50 were calculated using probit analysis.


### RESULTS

Table 1 shows the effect of different concentrations of EESP on *Aedes spp* larvae. The results showed that all larva were dead in T-treated group from the first observation time at 180 min which continued until 2880 min. In EESP-treated groups, the dead larvae were found at 1440 min and 2880 min observations ( $p < 0.05$ ). The number of dead larvae was increased that in line with the higher of EESP concentration. At 1440 min observation as follows CI(2.8 $\pm$ 1.7); CII(4 $\pm$ 1.8); CIII(4 $\pm$ 1.8); CIV(15.8 $\pm$ 1.5), and CV(17 $\pm$ 2.4), thus at 2880 min; CI(5 $\pm$ 1.4); CII(8 $\pm$ 1.4); CIII(13 $\pm$ 1.4); CIV(19 $\pm$ 1.4), and CV(21.3 $\pm$ 0.9). Contrarily, in W-treated group no larval mortality was found. These results revealed the biolarvicidal activity of EESP after 24 hours of exposure.

**Table:1** Effect of ethanolic extract of *Syzygium polyanthum* (Wight) (EESP) leaf on *Aedes spp* larvae instar III-IV mortality

Group	Number of larvae mortality by time (mean $\pm$ SD)			
	180'	360'	1440'	2880'
CI	0	0	2.8 $\pm$ 1.7	5 $\pm$ 1.4
CII	0	0	4 $\pm$ 1.8	8 $\pm$ 1.4
CIII	0	0	4 $\pm$ 1.8	13 $\pm$ 1.4
CIV	0	0	15.8 $\pm$ 1.5	19 $\pm$ 1.4
CV	0	0	17 $\pm$ 2.4	21.3 $\pm$ 0.9
T	25 $\pm$ 0	25 $\pm$ 0	25 $\pm$ 0	25 $\pm$ 0
W	0	0	0	0

Figure 1 shows that the percentage of *Aedes spp* larvae mortality increased with increasing of EESP concentration ie CI (20%), CII (32%), CIII(52%), CIV(76%) and CV (85%).



**Figure: 1** Percentage of *Aedes spp* larvae mortality after 48 hrs treatment

The present study showed that EESP with a concentration of 213 ppm and a time of 2410 minutes able to cause 50% mortality in the larvae of *Aedes spp* mosquitos (Table 2).

**Table: 2** LC50 and LT50 of ethanolic extract of *Syzygium polyanthum* (Wight) (EESP) leaf on *Aedes spp* larvae instar III-IV

Sample	LC50 (ppm)	LT50 (minutes)
EESP	213	2410

### DISCUSSION

The present study used the third and fourth stage of *Aedes* mosquito larvae based on WHO standards. At these stages, the instar not only more resistant to physical and mechanical factors such as displacement, limited space for living in water, but also having enough time to turn into an adult mosquito<sup>6,11-12</sup>.

Based on resistance and safety issues of chemical products, the efforts were turned to discover new compound of natural products derived from plants in mosquitoes control. New botanical natural products are believed to have following properties ie. effective, environment-friendly, easily biodegradable, inexpensive, and readily available in many areas of the world, no ill effect on non-target organisms and have novel modes of action. More than 2000 plant species known to have bioactivities as insecticide<sup>13-15</sup>.

*S. polyanthum* (Wight) leaf contained of alkaloid, atsiri oil, flavonoid, steroid, triterpenoid and saponin<sup>9</sup>. These phytochemical compounds were potentially toxic to insect. Alkaloids were reported can affect protein kinases which play a role in signal transduction, cell and tissue development and acetylcholinesterase inhibition. These compounds may damage the midgut and gastric caecum of larvae so that the larvae die<sup>16-18</sup>. Temephos could inhibit the cholinesterase enzyme, leading to impaired nerve activities due to the accumulation of acetylcholine<sup>19</sup>. Thus, alkaloids, atsiri oil, saponin, and flavonoids were also reported may damage nervous and respiratory system of larvae.

LC50 of EESP in the present study showed much lower than methanol extract of *S. polyanthum* (213 vs 6576.68 ppm) as reported by Tinneke and Puput, 2015, while Dwijanti, 2017 showed that water extract of *S. polyanthum* showed larvicidal activity at dose 2.5- 55%. The solvent used for extraction process of those different extracts affect the content of active compounds that can kill larvae.

### CONCLUSION

The present study showed that *Syzygium polyanthum* (Wight) leaf ethanolic extract have larvicidal activity on *Aedes spp* instar III-IV.

### CONFLICT OF INTEREST

The authors declare no conflict of interest.

### REFERENCES

- Walker TJPH, Johnson PH, Moreira LA, Iturbe-Ormaetxe I, Frentiu FD, McMeniman CJ, Lloyd AL. The wMel Wolbachia strain blocks dengue and invades caged *Aedes aegypti* populations. *Nature* 2011; 476(7361):450.

2. Owino EA. *Aedes spp* mosquitoes and emerging neglected diseases of Kenya. *Int J Mosq Res* 2018; 5(5):1-11.
3. Gizman, MG, Kouri G. Dengue: an update. *Lancet Infect Dis* 2002; 2:33-42.
4. Halstead SB. Dengue. *Lancet* 2007; 370:1644-1652.
5. Suaya JA, Shepard DS, Siqueira JB, Martelli CT, Lum LC, Tan LH, Armein B. Cost of dengue cases in eight countries in the Americas and Asia: a prospective study. *Am J Trop Med Hyg* 2009; 80(5):846-855.
6. Ardianto T. Pengaruh ekstrak bunga cengkeh (*Syzygium aromaticum* L.) terhadap mortalitas larva *Aedes aegypti* L. Surakarta: Fakultas Kedokteran Universitas Sebelas Maret 2008.
7. Kamaraj C, Rahuman AA, Bagavan A.. Antifeedant and larvicidal effects of plant extracts against *Spodoptera litura* (F.), *Aedes aegypti* L. and *Culex quinquefasciatus* Say. *Parasitol Res* 2008; 103(2):325-331.
8. Laurence M, Catherine D, Thomas G, Francis S, Marie V, Anna-Bella F. Phylogeography of *Aedes* (*Stegomyia*) *aegypti* (L.) and *Aedes* (*Stegomyia*) *albopictus* (Skuse) (Diptera: Culicidae) based on mitochondrial DNA variations. *Genetics Res* 2005; 86(1):1-11.
9. Widyawati T, Yusoff N, Asmawi M, Ahmad M. Antihyperglycemic effect of methanol extract of *Syzygium polyanthum* (Wight.) leaf in streptozotocin-induced diabetic rats. *Nutrients* 2015; 7(9):7764-7780.
10. Dwiyantri RD, Dediq R, Thuraidah A. Daya bunuh ekstrak air daun salam (*Syzygium polyathum*) terhadap larva *Aedes* sp. *Medi Lab Technol J* 2017; 3(1):17-21
11. World Health Organization. *Guidelines For Laboratory and Field Testing of Mosquito Larvicides*. World Health Organization 2005.
12. Tennyson S, Samraj DA, Jeyasundar D, Chalieu K. Larvicidal efficacy of plant oils against the dengue vector *Aedes aegypti* (L.) (diptera: culicidae). *Middle-East J Sci Res* 2013; 13(1):64-68.
13. Sukumar K, Perich MJ, Boobar LR. Botanical derivatives in mosquito control: A review. *J Am Mosq Control Asso* 1991; 7:210-237
14. Su T, Mulla MR. Oviposition bioassay responses of *Culex tarsalis* and *Culex quinquefasciatus* to neem products containing azadirachtin. *Entomol Exp Appl* 1999; 91:337-354
15. Sanei-Dehkordi A, Soleimani-Ahmadi M, Akbarzadeh K, Salim AY, Paksa A, Gorouhi MA, Mohammadi-Azni S. Chemical composition and mosquito larvicidal properties of essential oil from leaves of an Iranian indigenous plant *Zhumeria majdae*. *J Essent Oil Bear Pl* 2016; 19(6):1454-1461.
16. Ojha K, Pattabhiraiah M.. Evaluation of phytochemicals, larvicidal activity of *Jatropha curcas* seed oil against *Aedes aegypti*. *Int J Appl Res Stud* 2013; 2(12):1-12.
17. Ni'mah T, Oktarina R, Mahdalena V, Asyati D. Potensi ekstrak biji duku (*Lansium domesticum* Corr) terhadap *Aedes aegypti*. *Bul Penelit Kesehat* 2015; 43(2):131-136.
18. Velu K, Elumalai D, Hemalatha P, Babu M, Janaki A, Kaleena P. Phytochemical screening and larvicidal activity of peel extracts of *Arachis hypogaea* against chikungunya and malarial vectors. *Int J Mosq Res* 2015; 2(1):1-8.
19. Yulidar Y. Aktivitas gerak larva *Aedes aegypti* (Linn.) di bawah cekaman temefos. *Jurnal Edubio Tropika* 2014; 2(2).

